

Please ensure that your abstract fits into one column on one page and complies with the *Instructions to Authors* available from the Abstract Submission web page.

The Hawaiian PLUME Experiment and its Initial Data Assessment

G. LASKE^{1*}, A. MARKEE¹, J.A. ORCUTT¹, C.J. WOLFE²,
S.C. SOLOMON³, J.A. COLLINS⁴, R.S. DETRICK⁴, D.A.
BERCOVICI⁵ AND E.H. HAURI³

¹IGPP, SIO, UCSD, La Jolla, CA92093-0225

(*correspondence:glaske@ucsd.edu)

²University of Hawaii at Manoa, HI96822

³DTM, CIW, Washington, DC20015-1305

⁴WHOI, Woods Hole, MA02543-1050

⁵Yale University, New Haven, CT 06520-809

Hawaii has long been viewed as the textbook example for a plume-fed hotspot. Yet, the plume model has been contested because the collected seismic data to support or disprove it have so far been inconclusive. Compelling constraints on even the most basic features such as the plume conduit and its head have been elusive. One major problem that seismology has faced has been its complete reliance on land-based stations.

During the Jan. 2005 – May 2007 Hawaiian PLUME experiment we occupied nearly 70 sites with broad-band ocean bottom seismometers (OBSs). We collected continuous time series at a 1000 km-wide array that was augmented by 10 dedicated sites on the Hawaiian islands. The seismic data will facilitate the construction of body wave tomographic images of never-before obtained coverage and a depth-extent that reaches well into the lower mantle. The data will also be used to constrain the topography of mantle discontinuities through receiver functions and anisotropy through shear-wave splitting.

Here, we concentrate on the analyze of surface waves for which the used sensors provide ideal conditions for excellent depth resolution. For the first deployment phase, a 35-OBS array in Jan. 05 – Jan. 06, we analyzed data from upward of 95 suitable earthquakes and obtained internally consistent dispersion data. These allow us to image the lithosphere and upper asthenosphere. We find significant heterogeneity across the array. Higher phase velocities to the southeast of the island of Hawaii resemble those of 90 Myr old Pacific lithosphere. The lowest phase velocities are found to the west and northwest of the island of Hawaii. Our initial assessment is consistent the findings from the 1997/98 PELENET and SWELL pilot experiments that imaged a pronounced anomaly downstream from the island of Hawaii signifying profound changes to the 90 Myr old lithosphere, and possibly the underlying asthenosphere as well.